

GENERALIZED RENEWAL SEQUENCES AND INFINITELY DIVISIBLE LATTICE DISTRIBUTIONS

K. van HARN and F.W. STEUTEL

Department of Mathematics, Eindhoven University of Technology, Eindhoven, The Netherlands

Received December 1, 1975

We introduce an increasing set of classes Γ_α ($0 \leq \alpha \leq 1$) of infinitely divisible (i.d.) distributions on $\{0, 1, 2, \dots\}$, such that Γ_0 is the set of all compound-geometric distributions and Γ_1 the set of all compound-Poisson distributions, i.e. the set of all i.d. distributions on the non-negative integers. These classes are defined by recursion relations similar to those introduced by Katti [4] for Γ_1 and by Steutel [7] for Γ_0 . These relations can be regarded as generalizations of those defining the so-called renewal sequences (cf. [5] and [2]). Several properties of i.d. distributions now appear as special cases of properties of the Γ_α .

Infinite divisibility Lattice distributions Renewal sequences.

1. Introduction, summary and conventions

In [4] Katti gives the following theorem (see [8] for a simple proof).

Theorem 1.1. *A distribution $\{p_n\}_0^\infty$ on the non-negative integers, with $p_0 > 0$, is i.d. (i.e. compound-Poisson) iff there exist $r_n \geq 0$ ($n = 0, 1, 2, \dots$) such that*

$$(n+1)p_{n+1} = \sum_{k=0}^n p_k r_{n-k} \quad (n = 0, 1, 2, \dots); \quad (1.1)$$

the r_n then satisfy $\sum_0^\infty r_n / (n+1) < \infty$.

In [7] the following similar theorem is proved.

Theorem 1.2. *A distribution $\{p_n\}_0^\infty$ on the non-negative integers, with $p_0 > 0$, is compound-geometric iff there exist $r'_n \geq 0$ ($n = 0, 1, 2, \dots$) such that*

$$p_{n+1} = \sum_{k=0}^n p_k r'_{n-k} \quad (n = 0, 1, 2, \dots); \quad (1.2)$$

the r'_n then satisfy $\sum_0^\infty r'_n < 1$.

In fact, for any given distribution $\{p_n\}$ the equations (1.1) and (1.2) can be uniquely solved for r_n and r'_n (by Cramer's rule) to obtain explicit criteria for $\{p_n\}$ to be compound-Poisson, that is i.d., or compound-geometric. Conversely, any

sequence of non-negative r_n or r'_n , satisfying $\sum r_n/(n+1) < \infty$ or $\sum r'_n < 1$, by (1.1) or (1.2) uniquely determines a distribution $\{p_n\}$ that is compound-Poisson or compound-geometric. The equations (1.2) have the same structure as the equations defining the so-called renewal sequences (cf. [5]).

In this paper we consider a set of classes of i.d. distributions, that contains the classes defined by the Theorems 1.1 and 1.2 as special cases. For $0 \leq \alpha \leq 1$ we define the class Γ_α as follows.

Definition 1.3. A distribution $\{p_n\}_0^\infty$ on the non-negative integers, with $p_0 > 0$, belongs to Γ_α iff there exist $r_n(\alpha) \geq 0$ ($n = 0, 1, 2, \dots$) such that

$$c_n(\alpha)p_{n+1} = \sum_{k=0}^n p_k r_{n-k}(\alpha) \quad (n = 0, 1, 2, \dots); \quad (1.3)$$

the $r_n(\alpha)$ then satisfy

$$\sum_0^\infty r_n(\alpha)/c_n(\alpha) < \infty. \quad (1.4)$$

The quantities $c_n(\alpha)$, which will be specified in Section 2, are supposed to have the following properties.

$$\begin{aligned} c_n(0) &= 1, \quad c_n(1) = n + 1 \quad (n = 0, 1, 2, \dots) \\ c_n(\alpha) &\text{ is non-decreasing in both } n \text{ and } \alpha. \end{aligned} \quad (1.5)$$

Remark. The concluding statement of Definition 1.3 (see also Theorem 1.1) can be proved as follows:

$$1 - p_0 = \sum_{n=0}^\infty p_{n+1} = \sum_{k=0}^\infty p_k \sum_{n=k}^\infty r_{n-k}(\alpha)/c_n(\alpha) \geq p_0 \sum_{n=0}^\infty r_n(\alpha)/c_n(\alpha).$$

If $c(\alpha) := \lim_{n \rightarrow \infty} c_n(\alpha)$ is finite, then (see also Theorem 1.2)

$$\sum_0^\infty r_n(\alpha) < c(\alpha). \quad (1.6)$$

For given $c_n(\alpha)$ the sequences $\{p_n\}$ and $\{r_n(\alpha)\}$, satisfying (1.4) or (1.6), determine each other uniquely by means of (1.3). Clearly, Γ_0 is the class of compound-geometric distributions and Γ_1 the class of all i.d. distributions. It is well-known (see e.g. [6, p. 320] and [2]) that

$$\Gamma_0 \subset \Gamma_1. \quad (1.7)$$

In order to obtain a classification of the i.d. distributions on the non-negative integers, we wish to choose the $c_n(\alpha)$ such that

$$\Gamma_\alpha \subset \Gamma_\beta \quad \text{if } \alpha < \beta \quad (0 \leq \alpha < 1; 0 < \beta \leq 1), \quad (1.8)$$

and we shall say that in this case the $c_n(\alpha)$ satisfy (1.8). The main result of this paper is that this can be done, and, in fact, that the choice

$$c_n(\alpha) = \frac{1 - \alpha^{n+1}}{1 - \alpha} \quad (0 \leq \alpha \leq 1; n = 0, 1, 2, \dots), \quad (1.9)$$

with, of course, $c_n(1) = n + 1$, produces classes Γ_α satisfying (1.8). We prove several properties of the Γ_α , some of which are generalizations of well-known properties of i.d. distributions. We exhibit examples of distributions $\{p_n\}$ such that $\{p_n\} \in \Gamma_\alpha$ for $\alpha \geq \alpha_0$ and $\{p_n\} \notin \Gamma_\alpha$ for $\alpha < \alpha_0$, and we prove that $\bigcup_{\alpha < 1} \Gamma_\alpha$ is dense in Γ_1 .

We shall frequently use (probability) generating functions ((p.)g.f.'s), and we adopt the following notational conventions. The g.f.'s of $\{a_n\}$, $\{b_n\}$, ... are denoted by $A(z)$, $B(z)$, ..., those of $\{r_n(\alpha)\}$ and $\{c_n(\alpha)\}$ by $R_\alpha(z)$ and $C_\alpha(z)$. Clearly, $P(z)$ and $C_\alpha(z)$ converge for $|z| < 1$, whereas $R_\alpha(z)$ has a positive radius of convergence for every choice of the p_n , and is convergent for $|z| < 1$ if $\{p_n\} \in \Gamma_\alpha$, which will also be denoted by $P \in \Gamma_\alpha$.

As we consider only distributions on the non-negative integers with positive mass at 0, we use the term distribution in that sense.

For full proofs, details, examples and counterexamples we refer to [3]. For general information on i.d. distributions we refer to [1, chapter XII] and [6, Chapter 5].

2. The choice of $c_n(\alpha)$

Clearly, there are many sequences $\{c_n(\alpha)\}$ satisfying (1.5). The most obvious candidates, however, do not satisfy (1.8). We mention four of those, which, to avoid confusion, we denote by $c_n^*(\alpha)$. The following simple choices violate (1.8), as may be seen from counterexamples (cf. [3]).

$$c_n^*(\alpha) = 1 + \alpha n,$$

$$c_n^*(\alpha) = (n + 1)^\alpha,$$

$$c_n^*(\alpha) = (1 + \alpha n)^\alpha.$$

Rather more sophisticated, one may try to interpolate between the g.f.'s $C_0(z) = (1 - z)^{-1}$ and $C_1(z) = (1 - z)^{-2}$, e.g. by $C_\alpha^*(z) = (1 - z)^{-1-\alpha}$. This yields

$$c_n^*(\alpha) = \binom{\alpha + n}{n},$$

which somehow seems an attractive choice, but, unfortunately, also violates (1.8). We finally tried

$$C_\alpha(z) = (1 - z)^{-1}(1 - \alpha z)^{-1}, \quad (2.1)$$

corresponding to

$$c_n(\alpha) = \frac{1 - \alpha^{n+1}}{1 - \alpha} \quad (0 \leq \alpha \leq 1; n = 0, 1, 2, \dots), \quad (2.2)$$

with $c_n(1) = n + 1$. We shall prove in Section 3 that the $c_n(\alpha)$ given by (2.2) satisfy (1.8) (we note in passing that $\{c_n^*(\alpha)\}$ with $C_\alpha^*(z) = (1 - z)^{-1}(1 - \alpha z)^{-\alpha}$ does not).

The classes Γ_α as defined by (1.3) are now determined by the condition that the $r_n(\alpha)$, for a given distribution $\{p_n\}$ defined by

$$\frac{1 - \alpha^{n+1}}{1 - \alpha} p_{n+1} = \sum_{k=0}^n p_k r_{n-k}(\alpha), \quad (2.3)$$

should be non-negative, in which case $\sum_0^\infty r_n(\alpha) < (1 - \alpha)^{-1}$ if $\alpha < 1$ (cf. (1.6)). Equivalently, taking g.f.'s, R_α determined by

$$\frac{P(z) - P(\alpha z)}{z(1 - \alpha)} = P(z)R_\alpha(z) \quad (0 \leq \alpha < 1), \quad (2.4)$$

should be absolutely monotone (a.m.). We state this as a lemma.

Lemma 2.1. $P \in \Gamma_\alpha$ iff R_α is a.m., i.e. iff $1 - P(\alpha z)/P(z)$ is a.m. ($0 \leq \alpha < 1$).

If $\alpha = 0$ or $\alpha \uparrow 1$, we can solve (2.4) for $P(z)$, and obtain the well-known representations for Γ_0 and Γ_1 (see [1, p. 290] and [7, p. 84]).

Lemma 2.2. $P \in \Gamma_0$ iff P is of the form

$$P(z) = \frac{1 - p}{1 - pQ(z)}, \quad (2.5)$$

with $0 \leq p < 1$, and where Q is a p.g.f. with $Q(0) = 0$.

Lemma 2.3. $P \in \Gamma_1$ iff P is of the form

$$P(z) = \exp[\lambda\{Q(z) - 1\}], \quad (2.6)$$

with $\lambda > 0$, and where Q is a p.g.f. with $Q(0) = 0$.

3. The monotonicity of Γ_α

As a first step we prove that

$$\Gamma_\alpha \subset \Gamma_1 \quad (0 \leq \alpha < 1). \quad (3.1)$$

Theorem 3.1. If $\{p_n\} \in \Gamma_\alpha$, then $\{p_n\}$ is i.d.

Proof. From (2.4) it follows that

$$P(z) = \frac{P(\alpha z)}{1 - (1 - \alpha)zR_\alpha(z)}, \quad (3.2)$$

and iterating this equation we obtain

$$P(z) = \prod_{k=0}^{\infty} \frac{1 - \pi_k}{1 - \pi_k Q_k(z)}, \quad (3.3)$$

with $\pi_k = (1 - \alpha)\alpha^k R_\alpha(\alpha^k)$ and $Q_k(z) = zR_\alpha(\alpha^k z)/R_\alpha(\alpha^k)$. It is easily verified that the product in (3.3) is absolutely convergent for $|z| \leq 1$. Therefore, P is the limit of a sequence of products of p.g.f.'s in $\Gamma_0 \subset \Gamma_1$ (see (1.7)), and hence $P \in \Gamma_1$.

We state the analogue of Lemmas 2.2 and 2.3 as a corollary.

Corollary 3.2. *Let $0 \leq \alpha < 1$. Then $P \in \Gamma_\alpha$ iff P is of the form*

$$P(z) = \prod_{k=0}^{\infty} \frac{1 - pQ(\alpha^k)}{1 - pQ(\alpha^k z)}, \quad (3.4)$$

with $0 \leq p < 1$, and where Q is a p.g.f. with $Q(0) = 0$.

We now turn to the general theorem. The proof is fairly simple, but rather hard to find. We shall need the following lemma, which easily follows from Lemma 2.3.

Lemma 3.3. *If $P \in \Gamma_1$, then $P(z)/P(\gamma z)$ is a.m. for $0 \leq \gamma \leq 1$. In fact, $P(\gamma)P(z)/P(\gamma z)$ is a p.g.f. in Γ_1 .*

Theorem 3.4. $\Gamma_\alpha \subset \Gamma_\beta$ ($0 \leq \alpha < \beta \leq 1$).

Proof. We have to show (cf. Lemma 2.1) that, if $0 \leq \alpha < \beta < 1$ and if R_α is a.m., then R_β is a.m. To do so we rewrite (2.4) as

$$z(1 - \alpha)R_\alpha(z) = 1 - P(\alpha z)/P(z). \quad (3.5)$$

Now from (3.5) we subtract the same equality with z replaced by βz , and we obtain

$$z(1 - \alpha)\{R_\alpha(z) - \beta R_\alpha(\beta z)\} = \frac{P(\alpha\beta z)}{P(\beta z)} - \frac{P(\alpha z)}{P(z)},$$

or, equivalently,

$$\frac{z(1 - \alpha)}{P(\alpha z)}\{R_\alpha(z) - \beta R_\alpha(\beta z)\} = \frac{P(\alpha\beta z)}{P(\alpha z)P(\beta z)} - \frac{1}{P(z)}, \quad (3.6)$$

the right-hand side of which is symmetric in α and β . So, equating the left-hand side of (3.6) and the same expression with α and β interchanged, we get, after a simple rearrangement,

$$R_\beta(z) - \alpha R_\beta(\alpha z) = \frac{1 - \alpha}{1 - \beta} \frac{P(\beta z)}{P(\alpha z)} \{R_\alpha(z) - \beta R_\alpha(\beta z)\}. \quad (3.7)$$

By hypothesis R_α is a.m., and hence $R_\alpha(z) - \beta R_\alpha(\beta z)$, with coefficients $r_n(\alpha)(1 - \beta^{n+1})$, is a.m. By Lemma 3.3, and (3.1), $P(\beta z)/P(\alpha z)$ is a.m., and therefore $R_\beta(z) - \alpha R_\beta(\alpha z)$ is a.m., i.e. $r_n(\beta)(1 - \alpha^{n+1}) \geq 0$ ($n = 0, 1, 2, \dots$). It follows that $r_n(\beta) \geq 0$ ($n = 0, 1, 2, \dots$), and the theorem is proved.

Looking somewhat more precisely at the identity (3.7) one easily proves the following.

Corollary 3.5. *If for a given distribution R_α is a.m., and if $\alpha < \beta \leq 1$, then*

$$r_n(\beta) \geq \frac{1-\alpha}{1-\beta} \frac{1-\beta^{n+1}}{1-\alpha^{n+1}} r_n(\alpha) \quad (n = 0, 1, 2, \dots).$$

We restate Theorem 3.4 as a property that generalizes a property of renewal sequences proved in [2].

Corollary 3.6. *If for a given distribution $\{p_n\}$ the quantities $r_n(\alpha)$, for a fixed α ($0 \leq \alpha < 1$) defined by*

$$(1 + \alpha + \dots + \alpha^n) p_{n+1} = \sum_{k=0}^n p_k r_{n-k}(\alpha) \quad (n = 0, 1, 2, \dots), \quad (3.8)$$

are all non-negative, then also the quantities $r_n(\beta)$ defined by (3.8), with α replaced by β , are non-negative for all β with $\alpha < \beta \leq 1$.

Goldie's result in [2] is obtained by taking $\alpha = 0$ and $\beta = 1$.

4. Properties of Γ_α

The following theorem generalizes a well-known property of Γ_1 .

Theorem 4.1. *For $0 \leq \alpha \leq 1$ the class Γ_α is closed under weak convergence, i.e. if $P_n \in \Gamma_\alpha$ ($n = 1, 2, \dots$), and if $\lim_{n \rightarrow \infty} P_n(z) = P(z)$ is a p.g.f., then $P \in \Gamma_\alpha$.*

Proof. For $\alpha = 1$ the theorem is known (cf. [6]). From (2.4) it follows that if $P_n \rightarrow P$, then $R_{n,\alpha} \rightarrow R_\alpha$ ($0 \leq \alpha < 1$). As absolute monotonicity is preserved under point-wise convergence, R_α is a.m., and hence $P \in \Gamma_\alpha$ by Lemma 2.1.

Next we state some properties of Γ_α , which are well known, or trivial, when all Γ -classes are replaced by Γ_1 .

Theorem 4.2. *For $0 \leq \alpha < 1$,*

- (i) *if $P \in \Gamma_\alpha$, then $P(\gamma z)/P(\gamma) \in \Gamma_\alpha$ ($0 \leq \gamma \leq 1$);*
- (ii) *if $P \in \Gamma_\alpha$, then $P^\gamma \in \Gamma_\alpha$ ($0 \leq \gamma \leq 1$);*
- (iii) *if $P \in \Gamma_\alpha$, then*

$$P_{n,\gamma}(z) := \prod_{k=0}^{n-1} P(\gamma^k z)/P(\gamma^k) \in \Gamma_\gamma \quad (\alpha^{1/n} \leq \gamma \leq 1; n = 1, 2, \dots);$$

- (iv) *if $P \in \Gamma_\alpha$, then $P_\gamma(z) := P(\gamma)P(z)/P(\gamma z) \in \Gamma_\alpha$ ($0 \leq \gamma \leq 1$);*
- (v) *$P \in \Gamma_\alpha$ iff $P_\alpha(z) := P(\alpha)P(z)/P(\alpha z) \in \Gamma_0$.*

Proof. (i) follows directly from Lemma 2.1 and the fact that if $A(z)$ is a.m., then $A(\gamma z)$ is a.m.

(ii) The derivative of $1 - \{P(\alpha z)/P(z)\}^\gamma$ can be written as

$$\gamma \{P(z)/P(\alpha z)\}^{1-\gamma} \frac{d}{dz} \{1 - P(\alpha z)/P(z)\}. \quad (4.1)$$

By Lemma 3.3 we know that $P(\alpha)P(z)/P(\alpha z) \in \Gamma_1$ and hence $\{P(\alpha)P(z)/P(\alpha z)\}^{1-\gamma} \in \Gamma_1$, and hence is a.m. The second factor in (4.1) is a.m. by hypothesis, and (ii) follows by Lemma 2.1.

(iii) In order to use Lemma 2.1 again we calculate $1 - P_{n,\gamma}(\gamma z)/P_{n,\gamma}(z) = 1 - P(\gamma^n z)/P(z)$, which is a.m. by hypothesis for $\gamma^n = \alpha$, and hence by Theorem 3.4 for $\gamma^n \geq \alpha$.

(iv) We use Lemma 2.1 again, so we have to prove that

$$1 - P_\gamma(\alpha z)/P_\gamma(z) = \{P(\gamma z)/P(\alpha \gamma z)\}(1 - \alpha)z\{R_\alpha(z) - \gamma R_\alpha(\gamma z)\}$$

is a.m. This follows easily from Lemma 3.3 and the absolute monotonicity of R_α .

(v) We calculate $R_{0,\alpha}$ (i.e. R_0 corresponding to P_α), and get

$$R_{0,\alpha}(z) = (1 - \alpha)R_\alpha(z),$$

i.e. $R_{0,\alpha}$ is a.m. iff R_α is a.m. and so (v) follows.

We conclude this section by proving that $\bigcup_{0 \leq \alpha < 1} \Gamma_\alpha$ is dense in Γ_1 in the sense of weak convergence.

Theorem 4.3. *If $P \in \Gamma_1$, then there exists an increasing sequence $\{\alpha_n\}_1^\infty$, with $\alpha_n \rightarrow 1$, and a sequence of p.g.f.'s $P_n \in \Gamma_{\alpha_n}$ ($n = 1, 2, \dots$), such that*

$$P(z) = \lim_{n \rightarrow \infty} P_n(z) \quad (|z| \leq 1). \quad (4.2)$$

Proof. We prove the theorem for $0 \leq z \leq 1$; it is then easy to extend the result to $|z| \leq 1$. As (4.2) is trivial for $z = 1$, we take $z \in [0, 1)$. Now, let $P \in \Gamma_1$, then by Lemma 2.3

$$P(z) = \exp[\lambda\{Q(z) - 1\}]. \quad (4.3)$$

Take $\alpha_n = 1 - 1/n^2$, and for $n > \lambda$ define P_n by

$$P_n(z) = \prod_{k=0}^{n-1} \frac{1 - \lambda/n Q(\alpha_n^k)}{1 - \lambda/n Q(\alpha_n^k z)}.$$

By Theorem 4.2 (iii) we know that $P_n \in \Gamma_{\alpha_n}$. We rewrite P_n as

$$P_n(z) = \prod_{k=0}^{n-1} \left[1 + \frac{\lambda\{Q(z) - 1\}}{n} + \varepsilon_k(n) \right],$$

where it is not difficult (cf. [3] for details) to prove that $\varepsilon_k(n) = o(1/n)$ uniformly in k . It follows that $P_n(z) \rightarrow P(z)$ as given by (4.3).

5. Examples

We list a number of examples. The proofs of the following statements are simple, and are therefore omitted (cf. [3]).

$$1. \quad \prod_{k=0}^{n-1} \frac{1-p\alpha^k}{1-p\alpha^k z} \in \Gamma_\alpha \quad (0 \leq \alpha \leq 1; n = 1, 2, \dots).$$

$$2. \quad \frac{1-p}{1-pz} \frac{1-p\gamma}{1-p\gamma z} \in \Gamma_\alpha \text{ iff } \alpha \geq \gamma \quad (0 \leq p < 1; 0 \leq \gamma \leq 1).$$

$$3. \quad \left(\frac{1-p}{1-pz} \right)^{1+\varepsilon} \notin \Gamma_\alpha \quad (\varepsilon > 0; 0 < p < 1; 0 \leq \alpha < 1).$$

$$4. \quad \frac{1-p}{1-pz} \frac{1-p\gamma_1}{1-p\gamma_1 z} \frac{1-p\gamma_1\gamma_2}{1-p\gamma_1\gamma_2 z} \in \Gamma_\alpha \quad (0 \leq \gamma_1 \leq 1; 0 \leq \gamma_2 \leq 1; \alpha \geq \max(\gamma_1, \gamma_2); 0 \leq p < 1).$$

$$5. \quad \exp\{\lambda(z-1)\} \notin \Gamma_\alpha \quad (\lambda > 0; 0 \leq \alpha < 1).$$

6. If $P(z)$ is such that $R_{\alpha_0}(z) \equiv 1$, then $P \in \Gamma_\alpha$ iff $\alpha \geq \alpha_0$. Distributions of this type have been used in [3] to construct counterexamples to condition (1.8).

7. Let $p_0 = a/(1+a)$ and $p_n = \theta^n / \{(1+a)bn\}$ for $n \geq 1$, with $a > 0$, $b = -\log(1-\theta)$ and $0 < \theta < 1$. Then

$$\{p_n\} \in \Gamma_\alpha \text{ iff } ab \geq \frac{2}{1+\alpha} \quad (0 \leq \alpha \leq 1).$$

This can be proved by induction using (2.3). This example has also been considered in [4] and in [7], where it is noted that p_n is log-convex for $ab \geq 2$.

6. Concluding remarks

A more trivial classification of Γ_1 , with properties similar to those of Γ_α , is obtained by considering the classes Λ_u defined for $0 \leq u < \infty$ as follows:

$$P \in \Lambda_u \text{ iff } P(z) = \left\{ \frac{1-p}{1-pQ(z)} \right\}^u,$$

for some p with $0 \leq p < 1$ and some p.g.f. Q with $Q(0) = 0$.

Most of the properties of the Γ_α can be extended to similar classes of general (i.e. not necessarily lattice) distributions on $[0, \infty)$. Results of this kind will be published elsewhere.

It would be interesting to know whether other choices for $c_n(\alpha)$ exist, satisfying (1.5) and (1.8), especially whether there are such that also satisfy $c_n(\alpha) \sim n^\alpha$ ($n \rightarrow \infty$).

The solutions of the equations (2.3), with given $r_n(\alpha)$, non-negative and satisfying (1.4) or (1.6), may be regarded as generalized renewal sequences. It may be possible to extend some of the properties of renewal sequences proved in [5] to these more general sequences.

References

- [1] W. Feller, *An Introduction to Probability Theory and its Applications*, volume 1, 3rd ed. (Wiley, New York, 1968).
- [2] C.M. Goldie, A class of infinitely divisible distributions, *Proc. Cambridge Phil. Soc.* 63 (1967) 1141–1143.
- [3] K. van Harn and F.W. Steutel, Interpolating between the compound-geometric and compound-Poisson distributions, *Memorandum COSOR 75–10* (1975) (Department of Mathematics, Eindhoven University of Technology, Eindhoven, The Netherlands).
- [4] S.K. Katti, Infinite divisibility of integer valued random variables, *Ann. Math. Statist.* 38 (1967) 1306–1308.
- [5] D.G. Kendall, Renewal sequences and their arithmetic, *Symposium on Probability Methods in Analysis, Lecture Notes in Mathematics Nr. 31* (Springer, Berlin, Heidelberg, New York, 1967).
- [6] E. Lukács, *Characteristic Functions*, 2nd ed. (Griffin, London, 1970).
- [7] F.W. Steutel, Preservation of infinite divisibility under mixing, and related topics, *Math. Centre Tracts 33* (Math. Centre, Amsterdam, 1970).
- [8] F.W. Steutel, On the zeros of infinitely divisible densities, *Ann. Math. Statist.* 42 (1971) 812–815.